

APPLICATION UNDER UNITED STATES PATENT LAWS

Atty. Dkt. No. PW 277862
(M#)

Invention: NUCLEOTIDE SEQUENCES ENCODING THE dapC GENE AND PROCESS FOR THE PRODUCTION OF L-LYSINE

Inventor (s): Bettina MÖCKEL
Anke WEISSENBORN
Walter PFEFFERLE
Michael HARTMANN
Jörn KALINOWSKI
Alfred PÜHLER

Pillsbury Winthrop LLP
Intellectual Property Group
1100 New York Avenue, NW
Ninth Floor
Washington, DC 20005-3918
Attorneys
Telephone: (202) 861-3000

This is a:

- ☐ Provisional Application
- ☒ Regular Utility Application
- ☐ Continuing Application
 - ☒ The contents of the parent are incorporated by reference
- ☐ PCT National Phase Application
- ☐ Design Application
- ☐ Reissue Application
- ☐ Plant Application
- ☐ Substitute Specification
 - Sub. Spec. Filed _____
 - in App. No. _____ / _____
- ☐ Marked up Specification re
 - Sub. Spec. filed _____
 - In App. No. _____ / _____

SPECIFICATION

Nucleotide Sequences Encoding the dapC Gene and Process for the Production of L-Lysine

This application claims priority from German Application
No. 100 14 546.9, filed on March 23, 2000, the subject
5 matter of which is hereby incorporated herein by reference.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The invention provides nucleotide sequences encoding the
10 dapC gene and a process for the fermentative production of
L-lysine, using coryneform bacteria in which the dapC gene
(N-succinylaminoketopimelate transaminase gene) is
enhanced, in particular over-expressed.

2. Background Information

15 Amino acids, in particular L-lysine, are used in human
medicine and in the pharmaceuticals industry, but in
particular in animal nutrition.

It is known that amino acids are produced by fermentation
of strains of coryneform bacteria, in particular
20 Corynebacterium glutamicum. Due to their great
significance, efforts are constantly being made to improve
the production process. Improvements to the process may
relate to measures concerning fermentation technology, for
example stirring and oxygen supply, or to the composition
25 of the nutrient media, such as for example sugar
concentration during fermentation, or to working up of the
product by, for example, ion exchange chromatography, or to
the intrinsic performance characteristics of the
microorganism itself.

30 The performance characteristics of these microorganisms are
improved using methods of mutagenesis, selection and mutant
selection. In this manner, strains are obtained which are
resistant to antimetabolites, such as for example the
lysine analogue S-(2-aminoethyl)cysteine, or are

auxotrophic for regulatorily significant metabolites and produce L-amino acids, such as for example L-lysine.

For some years, methods of recombinant DNA technology have likewise been used to improve strains of *Corynebacterium* 5 which produce amino acids by amplifying individual amino acid biosynthesis genes and investigating the effect on amino acid production. Review articles on this subject may be found inter alia in Kinoshita ("Glutamic Acid Bacteria", in: *Biology of Industrial Microorganisms*, Demain and 10 Solomon (Eds.), Benjamin Cummings, London, UK, 1985, 115-142), Hilliger (*BioTec* 2, 40-44 (1991)), Eggeling (*Amino Acids* 6:261-272 (1994)), Jetten and Sinskey (*Critical Reviews in Biotechnology* 15, 73-103 (1995)) and Sahm et al. (15 (1996)).

SUMMARY OF THE INVENTION

It is an object of the invention to provide novel methods for the improved fermentative production of L-lysine.

Description of the Invention

20 L-lysine is used in human medicine, in the pharmaceuticals industry and in particular in animal nutrition. There is accordingly general interest in providing novel improved processes for the production of L-lysine.

Any subsequent mention of L-lysine or lysine should be 25 taken to mean not only the base, but also salts, such as for example lysine monohydrochloride or lysine sulfate.

The invention provides an isolated polynucleotide from coryneform bacteria containing at least one polynucleotide sequence selected from the group

30 a) polynucleotide which is at least 70% identical to a polynucleotide which encodes a polypeptide containing the amino acid sequence of SEQ ID no. 2,

- b) polynucleotide which encodes a polypeptide which contains an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID no. 2,
- c) polynucleotide which is complementary to the
5 polynucleotides of a) or b), or
- d) polynucleotide containing at least 15 successive nucleotides of the polynucleotide sequences of a), b) or c).

The invention also provides the polynucleotide according to
10 claim 1, wherein it preferably comprises replicable DNA containing:

- (i) the nucleotide sequence shown in SEQ ID no. 1, or
- (ii) at least one sequence which matches the sequence
15 (i) within the degeneration range of the genetic code, or
- (iii) at least one sequence which hybridizes with the complementary sequence to sequence (i) or (ii) and optionally
- (iv) functionally neutral sense mutations in (i).

20 The invention also provides

a polynucleotide according to claim 4, containing the nucleotide sequence as shown in SEQ ID no. 1,

a polynucleotide which encodes a polypeptide which contains the amino acid sequence as shown in SEQ ID no. 2,

25 a vector containing the polynucleotide according to claim 1, in particular a shuttle vector or the plasmid vector pXT-dapCexp, which is shown in Figure 2 and is deposited under number DSM 13254 in DSM 5715.

and coryneform bacteria acting as host cell which contain
30 the vector.

The invention also provides polynucleotides which substantially consist of a polynucleotide sequence, which are obtainable by screening by means of hybridization of a suitable gene library, which contains the complete gene
 5 having the polynucleotide sequence according to SEQ ID no. 1, with a probe which contains the sequence of the stated polynucleotide according to SEQ ID no. 1, or a fragment thereof, and isolation of the stated DNA sequence.

Polynucleotide sequences according to the invention are
 10 suitable as hybridization probes for RNA, cDNA and DNA in order to isolate full length cDNA which encode N-succinylaminoketopimelate transaminase and to isolate such cDNA or genes, the sequence of which exhibits a high level of similarity with that of the N-succinylaminoketopimelate
 15 transaminase gene.

Polynucleotide sequences according to the invention are furthermore suitable as primers for the production of DNA of genes which encode N-succinylaminoketopimelate transaminase by the polymerase chain reaction (PCR).

20 Such oligonucleotides acting as probes or primers contain at least 30, preferably at least 20, very particularly preferably at least 15 successive nucleotides. Oligonucleotides having a length of at least 40 or 50 nucleotides are also suitable.

25 "Isolated" means separated from its natural environment.

"Polynucleotide" generally relates to polyribonucleotides and polydeoxyribonucleotides, wherein the RNA or DNA may be unmodified or modified.

"Polypeptides" are taken to mean peptides or proteins which
 30 contain two or more amino acids connected by peptide bonds.

The polypeptides according to the invention include a polypeptide according to SEQ ID no. 2, in particular those having the biological activity of N-succinylaminoketopimelate transaminase and also those which are at least

70%, preferably at least 80%, identical to the polypeptide according to SEQ ID no. 2 and in particular are at least 90% to 95% identical to the polypeptide according to SEQ ID no. 2 and exhibit the stated activity.

- 5 The invention furthermore relates to a process for the fermentative production of amino acids, in particular L-lysine, using coryneform bacteria, which in particular already produce an amino acid and in which the nucleotide sequences which encode the dapC gene are enhanced, in
10 particular over-expressed.

In this connection, the term "enhancement" describes the increase in the intracellular activity of one or more enzymes in a microorganism, which enzymes are encoded by the corresponding DNA, for example by increasing the copy
15 number of the gene or genes, by using a strong promoter or a gene or allele which encodes a corresponding enzyme having elevated activity and optionally by combining these measures.

The microorganisms, provided by the present invention, may
20 produce L-amino acids, in particular L-lysine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. The microorganisms may comprise representatives of the coryneform bacteria in particular of the genus
25 *Corynebacterium*. Within the genus *Corynebacterium*, the species *Corynebacterium glutamicum* may in particular be mentioned, which is known in specialist circles for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in
30 particular of the species *Corynebacterium glutamicum*, are for example the known wild type strains

Corynebacterium glutamicum ATCC13032

Corynebacterium acetoglutamicum ATCC15806

Corynebacterium acetoacidophilum ATCC13870

35 *Corynebacterium thermoaminogenes* FERM BP-1539

Corynebacterium melassecola ATCC17965
 Brevibacterium flavum ATCC14067
 Brevibacterium lactofermentum ATCC13869 and
 Brevibacterium divaricatum ATCC14020

5 and L-lysine producing mutants or strains produced
 therefrom, such as for example

Corynebacterium glutamicum FERM-P 1709
 Brevibacterium flavum FERM-P 1708
 Brevibacterium lactofermentum FERM-P 1712
 10 Corynebacterium glutamicum FERM-P 6463
 Corynebacterium glutamicum FERM-P 6464
 Corynebacterium glutamicum DSM5715
 Corynebacterium glutamicum DSM12866 and
 Corynebacterium glutamicum DM58-1.

15 The inventors succeeded in isolating the novel dapC gene,
 which encodes the enzyme N-succinylaminoketopimelate
 transaminase (EC 2.6.1.17), from C. glutamicum.

The dapC gene, and also other genes from C. glutamicum, are
 isolated by initially constructing a gene library of this
 20 microorganism in E. coli. The construction of gene
 libraries is described in generally known textbooks and
 manuals. Examples which may be mentioned are the textbook
 by Winnacker, Gene und Klone, Eine Einführung in die
 Gentechnologie (Verlag Chemie, Weinheim, Germany, 1990) or
 25 the manual by Sambrook et al., Molecular Cloning, A
 Laboratory Manual (Cold Spring Harbor Laboratory Press,
 1989). One very well known gene library is that of E. coli
 K-12 strain W3110, which was constructed by Kohara et al.
 (Cell 50, 495-508 (1987)) in λ -vectors. Bathe et al.
 30 (Molecular and General Genetics, 252:255-265, 1996)
 describe a gene library of C. glutamicum ATCC13032, which
 was constructed using the cosmid vector SuperCos I (Wahl et
 al., 1987, Proceedings of the National Academy of Sciences
 USA, 84:2160-2164) in E. coli K-12 strain NM554 (Raleigh et
 35 al., 1988, Nucleic Acids Research 16:1563-1575). Börmann et
 al. (Molecular Microbiology 6(3), 317-326, 1992)) also

describe a gene library of *C. glutamicum* ATCC 13032, using cosmid pH79 (Hohn and Collins, Gene 11, 291-298 (1980)). A gene library of *C. glutamicum* in *E. coli* may also be produced using plasmids such as pBR322 (Bolivar, Life Sciences, 25, 807-818 (1979)) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are in particular those *E. coli* strains with restriction and recombination defects. One example of such a strain is the strain DH5 α mc^r, which has been described by Grant et al. (Proceedings of the National Academy of Sciences USA, 87 (1990) 4645-4649). The long DNA fragments cloned with the assistance of cosmids may then in turn be sub-cloned in usual vectors suitable for sequencing and then be sequenced, as described, for example, in Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977).

The novel DNA sequence from *C. glutamicum* which encodes the dapC gene and, as SEQ ID no. 1, is provided by the present invention, was obtained in this manner. The amino acid sequence of the corresponding protein was furthermore deduced from the above DNA sequence using the methods described above. SEQ ID no. 2 shows the resultant amino acid sequence of the product of the dapC gene.

Coding DNA sequences arising from SEQ ID no. 1 due to the degeneracy of the genetic code are also provided by the invention. DNA sequences which hybridize with SEQ ID no. 1 or parts of SEQ ID no. 1 are also provided by the invention. Conservative substitutions of amino acids in proteins, for example the substitution of glycine for alanine or of aspartic acid for glutamic acid, are known in specialist circles as "sense mutations", which result in no fundamental change in activity of the protein, i.e. they are functionally neutral. It is furthermore known that changes to the N and/or C terminus of a protein do not substantially impair or may even stabilize the function thereof. The person skilled in the art will find information in this connection inter alia in Ben-Bassat et

al. (Journal of Bacteriology 169:751-757 (1987)), in
 O'Regan et al. (Gene 77:237-251 (1989)), in Sahin-Toth et
 al. (Protein Sciences 3:240-247 (1994)), in Hochuli et al.
 (Bio/Technology 6:1321-1325 (1988)) and in known textbooks
 5 of genetics and molecular biology. Amino acid sequences
 arising in a corresponding manner from SEQ ID no. 2 are
 also provided by the invention.

Similarly, DNA sequences which hybridize with SEQ ID no. 1
 or portions of SEQ ID no. 1 are also provided by the
 10 invention. Finally, DNA sequences produced by the
 polymerase chain reaction (PCR) using primers obtained from
 SEQ ID no. 1 are also provided by the invention. Such
 oligonucleotides typically have a length of at least 15
 nucleotides.

15 The person skilled in the art may find instructions for
 identifying DNA sequences by means of hybridization inter
 alia in the manual "The DIG System Users Guide for Filter
 Hybridization" from Roche Diagnostics GmbH (Mannheim,
 Germany, 1993) and in Liebl et al. (International Journal
 20 of Systematic Bacteriology (1991) 41: 255-260). The person
 skilled in the art may find instructions for amplifying DNA
 sequences using the polymerase chain reaction (PCR) inter
 alia in the manual by Gait, Oligonucleotide Synthesis: A
 Practical Approach (IRL Press, Oxford, UK, 1984) and in
 25 Newton & Graham, PCR (Spektrum Akademischer Verlag,
 Heidelberg, Germany, 1994).

It has been found that coryneform bacteria produce L-lysine
 in an improved manner once the dapC gene has been over-
 expressed.

30 Over-expression may be achieved by increasing the copy
 number of the corresponding genes or by mutating the
 promoter and regulation region or the ribosome-binding site
 located upstream from the structural gene. Expression
 cassettes incorporated upstream from the structural gene
 35 act in the same manner. It is additionally possible to
 increase expression during fermentative L-lysine production

by means of inducible promoters. Expression is also improved by measures to extend the lifetime of the mRNA. Enzyme activity is moreover enhanced by preventing degradation of the enzyme protein. The genes or gene
 5 constructs may either be present in plasmids in a variable copy number or be integrated in the chromosome and amplified. Alternatively, over-expression of the genes concerned may also be achieved by modifying the composition of the media and culture conditions.

- 10 The person skilled in the art will find guidance in this connection inter alia in Martin et al. (Bio/Technology 3, 137-146 (1987)), in Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), in Eikmanns et al. (Gene 102, 93-98 (1991)), in EP
 15 0 472 869, in US 4,601,893, in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)), in Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), in LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), in WO 96/15246, in Malumbres et al.
 20 (Gene 134, 15-24 (1993)), in JP-A-10-229891, in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)), in Makrides (Microbiological Reviews 60:512-538 (1996)) and in known textbooks of genetics and molecular biology.

- 25 By way of example, the dapC gene according to the invention was over-expressed with the assistance of plasmids.

Suitable plasmids are those which are replicated in coryneform bacteria. Numerous known plasmid vectors, such as for example pZ1 (Menkel et al., Applied and
 30 Environmental Microbiology (1989) 64: 549-554), pEKEX1 (Eikmanns et al., Gene 102:93-98 (1991)) or pHS2-1 (Sonnen et al., Gene 107:69-74 (1991)) are based on the cryptic plasmids pHM1519, pBL1 or pGA1. Other plasmid vectors, such as for example those based on pCG4 (US-A 4,489,160), or
 35 pNG2 (Serwold-Davis et al., FEMS Microbiology Letters 66,

119-124 (1990)), or pAG1 (US-A 5,158,891) may be used in the same manner.

One example of a plasmid by means of which the dapC gene may be over-expressed is the E. coli-C. glutamicum shuttle
 5 vector pXT-dapCexp. The vector contains the replication region rep of plasmid pGA1, including the replication effector per (US-A- 5,175,108; Nesvera et al., Journal of Bacteriology 179, 1525-1532 (1997)), the tetA(Z) gene, which imparts tetracycline resistance, of plasmid pAG1 (US-
 10 A- 5,158,891; GenBank entry at the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA) with the accession number AF121000), together with the replication origin, the trc promoter, the termination regions T1 and T2 and the lacI^q gene (repressor of the lac
 15 operon of E. coli) of plasmid pTRC99A (Amann et al. (1988), Gene 69: 301-315).

The shuttle vector pXT-dapCexp is shown in Figure 2.

Further suitable plasmid vectors are those with the assistance of which gene amplification may be performed by
 20 integration into the chromosome, as has for example been described by Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)) for the duplication or amplification of the hom-thrB operon. In this method, the complete gene is cloned into a plasmid vector which can
 25 replicate in a host (typically E. coli), but not in C. glutamicum. Vectors which may be considered are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pGEM-T (Promega corporation, Madison, WI, USA),
 30 pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US-A 5,487,993), pCR®Blunt (Invitrogen, Groningen, Netherlands; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)) or pEM1 (Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516). The
 35 plasmid vector which contains the gene to be amplified is then transferred into the desired strain of C. glutamicum

by conjugation or transformation. The conjugation method is described, for example, in Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)).

Transformation methods are described, for example, in
5 Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous
10 recombination by means of "crossing over", the resultant strain contains at least two copies of the gene in question.

It has furthermore been found that, by replacing the amino acid L-proline in position 209 of the enzyme protein (c.f. SEQ ID no. 2) with another proteinogenic amino acid, in
15 particular L-leucine (c.f. SEQ ID no. 4), with the exception of L-proline, enhancement occurs and coryneform bacteria bearing the corresponding amino acid replacement produce L-lysine in an improved manner. The replacement of
20 L-proline with L-leucine in position 209 may preferably be achieved by replacing the nucleobase cytosine in position 716 with thymine as shown in SEQ ID no. 3.

Mutagenesis may be performed by conventional mutagenesis methods using mutagens such as for example N-methyl-N'-nitro-N-nitrosoguanidine or ultraviolet light. Mutagenesis
25 may also be performed by using in vitro methods such as for example treatment with hydroxylamine (Molecular and General Genetics 145, 101 pp (1978)) or mutagenic oligonucleotides (T.A. Brown: Gentechnologie für Einsteiger, Spektrum Akademischer Verlag, Heidelberg, 1993) or the polymerase
30 chain reaction (PCR), as is described in the manual by Newton and Graham (PCR, Spektrum Akademischer Verlag, Heidelberg, 1994).

The invention accordingly also provides DNA originating from coryneform bacteria which encodes
35 N-succinylaminoketopimelate transaminase, in which the amino acid sequence shown in SEQ ID no. 2 in position 209

is replaced with another amino acid, with the exception of L-proline. The invention also relates to coryneform bacteria which contain DNA in which the amino acid L-proline in position 209 of the enzyme protein (c.f. SEQ ID no. 2) is replaced with L-leucine (c.f. SEQ ID no. 4).

The invention furthermore provides coryneform bacteria which contain DNA in which the replacement of L-proline with L-leucine in position 209 proceeds by the replacement of the nucleobase cytosine in position 716 with thymine, as shown in SEQ ID no. 3.

It may additionally be advantageous for the production of L-lysine to amplify or over-express not only the dapC gene, but also one or more enzymes of the particular biosynthetic pathway, of glycolysis, of anaplerotic metabolism, or of amino acid export.

For the production of L-lysine, for example, it is thus possible in addition to the dapC gene simultaneously to enhance, in particular over-express or amplify, one or more genes selected from the group

- the lysC gene, which encodes a feed-back resistant aspartate kinase (Kalinowski et al. (1990), Molecular and General Genetics 224, 317-324),
- the asd gene, which encodes aspartate semialdehyde dehydrogenase (EP-A 0 219 027; Kalinowski et al. (1991), Molecular Microbiology 5:1197-1204, and Kalinowski et al. (1991), Molecular and General Genetics 224: 317-324),
- the dapA gene, which encodes dihydropicolinate synthase (EP-B 0 197 335),
- the dapB gene, which encodes dihydrodipicolinate reductase (GenBank entry accession number X67737; Pisabarro et al. (1993), Journal of Bacteriology, 175(9): 2743-2749),

- the dapD gene, which encodes tetrahydrodipicolinate succinylase (GenBank entry accession number AJ004934; Wehrmann et al. (1998), Journal of Bacteriology 180: 3159-3163),
- 5 • the dapE gene, which encodes N-succinyldiaminopimelate desuccinylase (GenBank entry accession number X81379; Wehrmann et al. (1994), Microbiology 140: 3349-3356),
- the dapF gene, which encodes diaminopimelate epimerase (DE: 199 43 587.1, DSM12968),
- 10 • the lysA gene, which encodes diaminopimelate decarboxylase (GenBank entry accession number X07563; Yeh et al. (1988), Molecular and General Genetics 212: 112-119),
- the ddh gene, which encodes diaminopimelate dehydrogenase
- 15 (Ishino et al. (1988), Agricultural and Biological Chemistry 52(11): 2903-2909),
- the lysE gene, which encodes lysine export (DE-A-195 48 222),
- the pyc gene, which encodes pyruvate carboxylase
- 20 (Eikmanns (1992), Journal of Bacteriology 174: 6076-6086),
- the mqo gene, which encodes malate:quinone oxidoreductase (Molenaar et al. (1998), European Journal of Biochemistry 254: 395-403),
- 25 • the zwal gene (DE: 19959328.0, DSM 13115),
- the gdh gene, which encodes glutamate dehydrogenase (Börmann et al. (1992), Molecular Microbiology 6, 317-326).

It is preferred simultaneously to enhance one or more genes
 30 selected from the group dapD, dapE and dapF.

It may furthermore be advantageous for the production of L-lysine, in addition to enhancing the dapC gene, optionally in combination with one or more genes selected from the group dapD, dapE and dapF, simultaneously to attenuate

- 5 • the pck gene, which encodes phosphoenolpyruvate carboxykinase (DE 199 50 409.1, DSM 13047) or
- the pgi gene, which encodes glucose 6-phosphate isomerase (US 09/396,478, DSM 12969), or
- the poxB gene, which encodes pyruvate oxidase
- 10 (DE: 19951975.7, DSM 13114), or
- the zwa2 gene (DE: 19959327.2, DSM 13113), or
- the sucC or sucD genes which encode succinyl CoA synthetase (DE: 19956686.0).

It may furthermore be advantageous for the production of L-lysine, in addition to enhancing the dapC gene, optionally

15 in combination with one or more genes selected from the group dapD, dapE and dapF, to suppress unwanted secondary reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Over-production of Microbial

20 Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

For the purposes of L-lysine production, the microorganisms produced according to the invention may be cultured continuously or discontinuously using the batch process or

25 the fed batch process or repeated fed batch process. A summary of known culture methods is given in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und

30 periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must adequately satisfy the requirements of the particular strains. Culture media for

various microorganisms are described in "Manual of Methods for General Bacteriology" from the American Society for Bacteriology (Washington D.C., USA, 1981).

Carbon sources which may be used are sugars and
 5 carbohydrates, such as glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose for example, oils and fats, such as soya oil, sunflower oil, peanut oil and coconut oil for example, fatty acids, such as palmitic acid, stearic acid and linoleic acid for example, alcohols,
 10 such as glycerol and ethanol for example, and organic acids, such as acetic acid for example. These substances may be used individually or as a mixture.

Nitrogen sources which may be used comprise organic compounds containing nitrogen, such as peptones, yeast
 15 extract, meat extract, malt extract, corn steep liquor, soya flour and urea or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate. The nitrogen sources may be used individually or as a mixture.

20 Phosphorus sources which may be used are phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding salts containing sodium. The culture medium has additionally to contain salts of metals, such as magnesium sulfate or iron sulfate for example,
 25 which are necessary for growth. Finally, essential growth-promoting substances such as amino acids and vitamins may also be used in addition to the above-stated substances. Suitable precursors may furthermore be added to the culture medium. The stated feed substances may be added to the
 30 culture as a single batch or be fed appropriately during culturing.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or ammonia water, or acidic compounds, such as phosphoric acid or sulfuric acid, are used
 35 appropriately to control the pH of the culture. Foaming may be controlled by using antifoaming agents such as fatty

acid polyglycol esters for example. Plasmid stability may be maintained by the addition to the medium of suitable selectively acting substances, for example antibiotics. Oxygen or oxygen-containing gas mixtures, such as air for example, are introduced into the culture in order to maintain aerobic conditions. The temperature of the culture is normally from 20°C to 45°C and preferably from 25°C to 40°C. The culture is continued until the maximum quantity of lysine has formed. This aim is normally achieved within 10 to 160 hours.

Analysis of L-lysine may be performed by anion exchange chromatography with subsequent ninhydrin derivation, as described in Spackman et al. (Analytical Chemistry, 30, (1958), 1190).

The following microorganism has been deposited with Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ, Braunschweig, Germany) in accordance with the Budapest Treaty:

- *Corynebacterium glutamicum* strain DSM5715/pXT-dapCexp as DSM 13254.

The process according to the invention serves in the fermentative production of L-lysine.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Map of plasmid pEC-XT99A

Figure 2: Map of plasmid pXT-dapCexp

The abbreviations and names are defined as follows.

per:	Gene for controlling copy number from pGA1
oriE:	Plasmid-coded replication origin of <i>E. coli</i>
rep:	Plasmid-coded replication origin from <i>C. glutamicum</i> plasmid pGA1
P _{trc} :	trc promoter from pTRC99A
T1, T2:	Terminator regions 1 and 2 from pTRC99A
lacIq:	Repressor gene of the Lac operon

Tet: Resistance gene for tetracycline
 dapC: dapC gene from *C. glutamicum*
 EcoRI: Restriction site of the restriction enzyme
 EcoRI
 5 EcoRV: Restriction site of the restriction enzyme
 EcoRV
 HindIII: Restriction site of the restriction enzyme
 HindIII
 KpnI: Restriction site of the restriction enzyme
 10 KpnI
 SalI: Restriction site of the restriction enzyme
 SalI
 SmaI: Restriction site of the restriction enzyme
 SmaI
 15 NdeI: Restriction site of the restriction enzyme
 NdeI
 BamHI: Restriction site of the restriction enzyme
 BamHI
 NcoI: Restriction site of the restriction enzyme
 20 NcoI
 XbaI: Restriction site of the restriction enzyme
 XbaI
 SacI: Restriction site of the restriction enzyme
 SacI

25

DETAILED DESCRIPTION OF THE INVENTION

The present invention is illustrated in greater detail by the following practical examples.

Example 1

30 Production of a genomic cosmid gene library from
Corynebacterium glutamicum ATCC 13032

Chromosomal DNA from *Corynebacterium glutamicum* ATCC 13032
 was isolated as described in Tauch et al., (1995, Plasmid
 33:168-179) and partially cleaved with the restriction
 35 enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany,
 product description Sau3AI, code no. 27-0913-02). The DNA

fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, product description SAP, code no. 1758250). The DNA of cosmid vector SuperCos1 (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164), purchased from Stratagene (La Jolla, USA, product description SuperCos1 Cosmid Vector Kit, code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, product description XbaI, code no. 27-0948-02) and also dephosphorylated with shrimp alkaline phosphatase. The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, product description BamHI, code no. 27-0868-04). Cosmid DNA treated in this manner was mixed with the treated ATCC 13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, product description T4 DNA Ligase, code no. 27-0870-04). The ligation mixture was then packed in phages using Gigapack II XL Packing Extracts (Stratagene, La Jolla, USA, product description Gigapack II XL Packing Extract, code no. 200217). E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Research 16:1563-1575) was infected by suspending the cells in 10 mM MgSO₄ and mixing them with an aliquot of the phage suspension. The cosmid library was infected and titred as described in Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) with 100 mg/l of ampicillin. After overnight incubation at 37°C, individual recombinant clones were selected.

Example 2

Isolation and sequencing of the dapC gene

Cosmid DNA from an individual colony was isolated in accordance with the manufacturer's instructions using the Qiaprep Spin Miniprep Kit (product no. 27106, Qiagen, Hilden, Germany) and partially cleaved with the restriction

enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, product description Sau3AI, product no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, product description SAP, product no. 1758250). Once separated by gel electrophoresis, the cosmid fragments of a size of 1500 to 2000 bp were isolated using the QiaExII Gel Extraction Kit (product no. 20021, Qiagen, Hilden, Germany). The DNA of the sequencing vector pZero-1 purchased from Invitrogen (Groningen, Netherlands, product description Zero Background Cloning Kit, product no. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, product description BamHI, product no. 27-0868-04). Ligation of the cosmid fragments into the sequencing vector pZero-1 was performed as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated into the E. coli strain DH5αMCR (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A., 87:4645-4649) (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7) and plated out onto LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l of Zeocin. Plasmids of the recombinant clones were prepared using the Biorobot 9600 (product no. 900200, Qiagen, Hilden, Germany). Sequencing was performed using the dideoxy chain termination method according to Sanger et al. (1977, Proceedings of the National Academies of Sciences U.S.A., 74:5463-5467) as modified by Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (product no. 403044, Weiterstadt, Germany) was used. Separation by gel electrophoresis and analysis of the sequencing reaction was performed in a "Rotiphorese NF" acrylamide/bisacrylamide gel (29:1) (product no. A124.1, Roth, Karlsruhe, Germany) using the "ABI Prism 377"

sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The resultant raw sequence data were then processed using the Staden software package (1986, Nucleic Acids Research, 14:217-231), version 97-0. The individual sequences of the pZero1 derivatives were assembled into a cohesive contig. Computer-aided coding range analysis was performed using XNIP software (Staden, 1986, Nucleic Acids Research, 14:217-231). Further analysis was performed using the "BLAST search programs" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402), against the non-redundant database of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).

The resultant nucleotide sequence of the dapC gene is stated in SEQ ID no. 1. Analysis of the nucleotide sequence revealed an open reading frame of 1101 base pairs, which was designated the dapC gene. The dapC gene encodes a polypeptide of 367 amino acids, which is shown in SEQ ID no. 2.

20 Example 3

Production of a shuttle vector pXT-dapCexp for enhancing the dapC gene in *C. glutamicum*

3.1. Cloning of the dapC gene

Chromosomal DNA was isolated from strain ATCC 13032 using the method of Eikmanns et al. (Microbiology 140: 1817-1828 (1994)). On the basis of the sequence of the dapC gene for *C. glutamicum* known from Example 2, the following oligonucleotides were selected for the polymerase chain reaction (c.f. also SEQ ID no. 5 and 6):

30 DapC (dCex1):

5` GAT CTA (GAA TTC) GCC TCA GGC ATA ATC TAA CG 3`

DapC (dCexna2):

5` GAT CTA (TCT AGA) CAG AGG ACA AGG CAA TCG GA 3`

The stated primers were synthesized by the company ARK Scientific GmbH Biosystems (Darmstadt, Germany) and the PCR reaction performed in accordance with the standard PCR method of Innis et al. (PCR Protocols. A Guide to Methods and Applications, 1990, Academic Press) using Pwo polymerase from Roche Diagnostics GmbH (Mannheim, Germany). By means of the polymerase chain reaction, the primers permit the amplification of an approx. 1.6 kb DNA fragment, which bears the dapC gene. Moreover, the primer DapC (dCex1) contains the sequence for the restriction site of the restriction endonuclease EcoRI, and the primer DapC (dCexna2) contains the restriction site of the restriction endonuclease XbaI, which are indicated between brackets in the above-stated nucleotide sequence.

The amplified approx. 1.6 kb DNA fragment, which bears the dapC gene, was ligated into the vector pCR®Blunt II (Bernard et al., Journal of Molecular Biology, 234:534-541 (1993)) using the Zero Blunt™ Kit from Invitrogen Corporation (Carlsbad, CA, USA; catalogue number K2700-20).

The E. coli strain Top10 was then transformed with the ligation batch in accordance with the kit manufacturer's instructions (Invitrogen Corporation, Carlsbad, CA, USA). Plasmid-bearing cells were selected by plating the transformation batch out onto LB agar (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) which had been supplemented with 25 mg/l of kanamycin. Plasmid DNA was isolated from a transformant using the QIAprep Spin Miniprep Kit from Qiagen (Hilden, Germany) and verified by restriction with the restriction enzymes XbaI and EcoRI and subsequent (0.8%) agarose gel electrophoresis. The DNA sequence of the amplified DNA fragment was verified by sequencing. The plasmid was named pCRdapC. The strain was designated E. coli Top10 / pCRdapC.

3.2. Production of the E. coli-C. glutamicum shuttle vector pEC-XT99A

The *E. coli* expression vector pTRC99A (Amann et al. 1988, Gene 69:301-315) was used as the starting vector for constructing the *E. coli*-*C. glutamicum* shuttle expression vector pEC-XT99A. After BspHI restriction cleavage (Roche
 5 Diagnostics GmbH, Mannheim, Germany, production description BspHI, product no. 1467123) and subsequent Klenow treatment (Amersham Pharmacia Biotech, Freiburg, Germany, product
 description Klenow Fragment of DNA Polymerase I, product
 no. 27-0928-01; method according to Sambrook et al., 1989,
 10 Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the ampicillin resistance gene (*bla*) was replaced by the tetracycline resistance gene of *C. glutamicum* plasmid pAG1 (GenBank accession no. AF121000). To this end, the region bearing the resistance gene was cloned as an
 15 AluI fragment (Amersham Pharmacia Biotech, Freiburg, Germany, product description AluI, product no. 27-0884-01) into the linearized *E. coli* expression vector pTRC99A. Ligation was performed as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring
 20 Harbor), the DNA mixture being incubated overnight with T4 ligase (Amersham Pharmacia Biotech, Freiburg, Germany, product description T4 DNA Ligase, product no. 27-0870-04). This ligation mixture was then electroporated into the *E. coli* strain DH5 α mc^r (Grant, 1990, Proceedings of the
 25 National Academy of Sciences U.S.A., 87:4645-4649) (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7). The constructed *E. coli* expression vector was designated pXT99A.

Plasmid pGA1 (Sonnen et al. 1991, Gene, 107:69-74) was used
 30 as the basis for cloning a minimal replicon from *Corynebacterium glutamicum*. By means of BalI/PstI restriction cleavage (Promega GmbH, Mannheim, Germany, production description BalI, product no. R6691; Amersham Pharmacia Biotech, Freiburg, Germany, production
 35 description PstI, product no. 27-0976-01) of the pGA1 vector, it proved possible to clone a 3484 bp fragment into the pK18mob2 vector (Tauch et al., 1998, Archives of Microbiology 169:303-312) which had been fragmented with

SmaI und PstI (Amersham Pharmacia Biotech, Freiburg, Germany, product description SmaI, product no. 27-0942-02, product description PstI, product no. 27-0976-01). An 839 bp fragment was deleted by BamHI/XhoI restriction cleavage

5 (Amersham Pharmacia Biotech, Freiburg, Germany, product description BamHI, product no. 27-086803, product description XhoI, product no. 27-0950-01) and subsequent Klenow treatment (Amersham Pharmacia Biotech, Freiburg, Germany, product description Klenow Fragment of DNA

10 Polymerase I, product no. 27-0928-01; method according to Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor). The *C. glutamicum* minimal replicon could be cloned into the *E. coli* expression vector pXT99A as a 2645 bp fragment from the construct which had

15 been religated with T4 ligase (Amersham Pharmacia Biotech, Freiburg, Germany, product description T4 DNA Ligase, product no. 27-0870-04). To this end, the DNA of the construct bearing the minimal replicon was cleaved with the restriction enzymes KpnI (Amersham Pharmacia Biotech,

20 Freiburg, Germany, product description KpnI, product no. 27-0908-01) and PstI (Amersham Pharmacia Biotech, Freiburg, Germany, product description PstI, product no. 27-0886-03) and then a 3'-5'-exonuclease treatment (Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring

25 Harbor) was performed by means of Klenow polymerase (Amersham Pharmacia Biotech, Freiburg, Germany, product description Klenow Fragment of DNA Polymerase I, product no. 27-0928-01).

In a parallel batch, the *E. coli* expression vector pXT99A

30 was cleaved with the restriction enzyme RsrII (Roche Diagnostics, Mannheim, Germany, product description RsrII, product no. 1292587) and prepared for ligation with Klenow polymerase (Amersham Pharmacia Biotech, Freiburg, Germany, Klenow Fragment of DNA Polymerase I, product no. 27-0928-

35 01). Ligation of the minimal replicon with the vector construct pXT99A was performed as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight

with T4 ligase (Amersham Pharmacia Biotech, Freiburg, Germany, product description T4 DNA Ligase, product no. 27-0870-40).

The E. coli-C. glutamicum shuttle expression vector pEC-XT99A constructed in this manner was transferred into C. glutamicum DSM5715 by electroporation (Liebl et al., 1989, FEMS Microbiology Letters, 53:299-303). Transformant selection proceeded on LBHIS agar consisting of 18.5 g/l of brain-heart infusion bouillon, 0.5 M sorbitol, 5 g/l of Bacto tryptone, 2.5 g/l of Bacto yeast extract, 5 g/l of NaCl and 18 g/l of Bacto agar, which had been supplemented with 5 mg/l of tetracycline. Incubation was performed for 2 days at 33°C.

Plasmid DNA was isolated from a transformant using the conventional methods (Peters-Wendisch et al., 1998, Microbiology, 144, 915 - 927), cut with the restriction endonuclease HindIII and the plasmid verified by subsequent agarose gel electrophoresis.

The resultant plasmid construct was named pEC-XT99A and is shown in Figure 1. The strain obtained by electroporation of plasmid pEC-XT99A into Corynebacterium glutamicum DSM5715 was named DSM5715/pEC-XT99A and has been deposited with Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ, Braunschweig, Germany) in accordance with the Budapest Treaty.

3.3. Cloning of dapC in the E. coli-C. glutamicum shuttle vector pEC-XT99A

The vector used was the E. coli-C. glutamicum shuttle vector pEC-XT99A described in Example 3.2. DNA from this plasmid was completely cleaved with the restriction enzymes EcoRI and XbaI and then dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, product description SAP, product no. 1758250).

The dapC gene was isolated from the plasmid pCRdapC described in Example 3.1. by complete cleavage with the enzymes EcoRI and XbaI. The approx. 1600 bp dapC fragment was isolated from the agarose gel using the QiaExII Gel
 5 Extraction Kit (product no. 20021, Qiagen, Hilden, Germany).

The dapC fragment obtained in this manner was mixed with the prepared pEC-XT99A vector and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany,
 10 product description T4 DNA Ligase, code no. 27-0870-04). The ligation batch was then transformed into E. coli strain DH5 α (Hanahan, in: DNA Cloning. A Practical Approach, Vol. I, IRL-Press, Oxford, Washington DC, USA). Plasmid-bearing cells were selected by plating the transformation batch out
 15 onto LB agar (Lennox, 1955, Virology, 1:190) with 5 mg/l of tetracycline. After overnight incubation at 37°C, individual recombinant clones were selected. Plasmid DNA was isolated from a transformant in accordance with the manufacturer's instructions using the Qiaprep Spin Miniprep
 20 Kit (product no. 27106, Qiagen, Hilden, Germany) and cleaved with the restriction enzymes EcoRI and XbaI in order to verify the plasmid by subsequent agarose gel electrophoresis. The resultant plasmid was named pXT-dapCexp. It is shown in Figure 2.

25 Example 4

Transformation of strain DSM5715 with plasmid pXT-dapCexp

Strain DSM5715 was then transformed with plasmid pXT-dapCexp using the electroporation method described by Liebl et al. (FEMS Microbiology Letters, 53:299-303 (1989)).
 30 Transformant selection proceeded on LBHIS agar consisting of 18.5 g/l of brain-heart infusion bouillon, 0.5 M sorbitol, 5 g/l of Bacto tryptone, 2.5 g/l of Bacto yeast extract, 5 g/l of NaCl and 18 g/l of Bacto agar, which had been supplemented with 5 mg/l of tetracycline. Incubation
 35 was performed for 2 days at 33°C.

Plasmid DNA was isolated from a transformant using the conventional methods (Peters-Wendisch et al., 1998, Microbiology, 144, 915 - 927), cut with the restriction endonucleases EcoRI and XbaI and the plasmid was verified
 5 by subsequent agarose gel electrophoresis. The resultant strain was named DSM5715/pXT-dapCexp and has been deposited as DSM 13254 with Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ, Braunschweig, Germany) in accordance with the Budapest Treaty.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
220

Example 5

Production of L-lysine

The *C. glutamicum* strain DSM5715/pXT-dapCexp obtained in Example 4 was cultured in a nutrient medium suitable for the production of lysine and the lysine content of the culture supernatant was determined.

To this end, the strain was initially incubated for 24 hours at 33°C on an agar plate with the appropriate antibiotic (brain/heart agar with tetracycline (5 mg/l)). Starting from this agar plate culture, a preculture was inoculated (10 ml of medium in a 100 ml Erlenmeyer flask). The complete medium CgIII was used as the medium for this preculture.

Medium Cg III

NaCl	2.5 g/l
Bacto peptone	10 g/l
Bacto yeast extract	10 g/l
Glucose (separately autoclaved)	2% (w/v)

The pH value was adjusted to pH 7.4.

15

Tetracycline (5 mg/l) was added to this medium. The preculture was incubated for 16 hours at 33°C on a shaker at 240 rpm. A main culture was inoculated from this preculture, such that the initial OD (660 nm) of the main culture was 0.1. Medium MM was used for the main culture.

Medium MM

CSL (Corn Steep Liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (separately autoclaved)	50 g/l
$(\text{NH}_4)_2\text{SO}_4$	25 g/l
KH_2PO_4	0.1 g/l
$\text{MgSO}_4 \cdot 7 \text{ H}_2\text{O}$	1.0 g/l
$\text{CaCl}_2 \cdot 2 \text{ H}_2\text{O}$	10 mg/l
$\text{FeSO}_4 \cdot 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-leucine (sterile-filtered)	0.1 g/l
CaCO_3	25 g/l

CSL, MOPS and the salt solution were adjusted to pH 7 with ammonia water and autoclaved. The sterile substrate and
 5 vitamin solutions, together with the dry-autoclaved CaCO_3 are then added.

Culturing is performed in a volume of 10 ml in a 100 ml Erlenmeyer flask with flow spoilers. Tetracycline (5 mg/l) was added. Culturing was performed at 33°C and 80%
 10 atmospheric humidity.

After 72 hours, the OD was determined at a measurement wavelength of 660 nm using a Biomek 1000 (Beckmann Instruments GmbH, Munich). The quantity of lysine formed

was determined using an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivation with ninhydrin detection.

Table 1 shows the result of the test.

5

Table 1

Strain	OD (660 nm)	Lysine HCl 25 g/l
DSM5715	7.0	13.7
DSM5715/pXT-dapCexp	7.1	14.7

SEQUENCE LISTING

<110> Degussa-Huels AG

5 <120> Nucleotide Sequences Coding for the dapC Gene and Process for
the Production of L-Lysine.

<130> 990217 BT

10 <140>

<141>

<160> 6

15 <170> PatentIn Ver. 2.1

<210> 1

<211> 1300

<212> DNA

20 <213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (91)..(1191)

25 <223> dapC-Gene

<400> 1

gggctcccca ggtggtgctg ctaagcttgg accacaagat tttgatcacc caatgatcgc 60

30 tgcgctgctg cctcaggcat aatctaacgc atg acc tct cgc acc ccg ctt gtt 114
Met Thr Ser Arg Thr Pro Leu Val
1 5

35 tct gtt ctt cct gat ttt ccg tgg gat tgc ctc gct tcc gca aaa gcc 162
Ser Val Leu Pro Asp Phe Pro Trp Asp Ser Leu Ala Ser Ala Lys Ala
10 15 20

40 aaa gct gcg tct cac ccg gat ggg atc gtg aat ctt tct gtt ggc act 210
Lys Ala Ala Ser His Pro Asp Gly Ile Val Asn Leu Ser Val Gly Thr
25 30 35 40

45 ccg gtt gat ccg gtc gcg ccc agc att cag atc gcg ttg gca gaa gca 258
Pro Val Asp Pro Val Ala Pro Ser Ile Gln Ile Ala Leu Ala Glu Ala
45 50 55

50 gcg ggg ttt tgc ggt tac cct caa acc atc ggc acc ccg gaa ctc cgc 306
Ala Gly Phe Ser Gly Tyr Pro Gln Thr Ile Gly Thr Pro Glu Leu Arg
60 65 70

55 gca gcc atc agg ggc gcg ctt gag cgg cgc tac aac atg aca aag ctt 354
Ala Ala Ile Arg Gly Ala Leu Glu Arg Arg Tyr Asn Met Thr Lys Leu
75 80 85

60 gtc gac gcc tcc ctc ctc ccc gtc gtg ggt acc aag gag gca att gcc 402
Val Asp Ala Ser Leu Leu Pro Val Val Gly Thr Lys Glu Ala Ile Ala
90 95 100

60 ctt ctt cca ttc gcg ttg ggt att tcc ggc acc gtt gtc atc cca gag 450
Leu Leu Pro Phe Ala Leu Gly Ile Ser Gly Thr Val Val Ile Pro Glu
105 110 115 120

att gcg tac cca acc tac gaa gtc gct gtc gtg gcc gca gga tgc acc 498
 Ile Ala Tyr Pro Thr Tyr Glu Val Ala Val Val Ala Ala Gly Cys Thr
 125 130 135

5 gtg ttg cgt tct gat tcg ctg ttt aag ctc ggc ccg cag atc ccg tcg 546
 Val Leu Arg Ser Asp Ser Leu Phe Lys Leu Gly Pro Gln Ile Pro Ser
 140 145 150

10 atg atg ttt atc aac tca cca tcc aac ccc aca ggc aag gtt ctg ggc 594
 Met Met Phe Ile Asn Ser Pro Ser Asn Pro Thr Gly Lys Val Leu Gly
 155 160 165

15 atc cca cac ttg cgc aag gtt gtg aag tgg gcg cag gaa aac aac gtg 642
 Ile Pro His Leu Arg Lys Val Val Lys Trp Ala Gln Glu Asn Asn Val
 170 175 180

20 atc ctc gca gct gat gaa tgc tac ttg ggt ctt ggc tgg gac gat gaa 690
 Ile Leu Ala Ala Asp Glu Cys Tyr Leu Gly Leu Gly Trp Asp Asp Glu
 185 190 195 200

aac cca ccg atc tca att ttg gat cca cgt gtc tgc gat ggc gac cac 738
 Asn Pro Pro Ile Ser Ile Leu Asp Pro Arg Val Cys Asp Gly Asp His
 205 210 215

25 acc aac ttg atc gcc att cac tcg ctg tct aaa acc tca aac ctc gct 786
 Thr Asn Leu Ile Ala Ile His Ser Leu Ser Lys Thr Ser Asn Leu Ala
 220 225 230

30 tct tac cgc gca ggt tac ctc gtt ggc gat cca gcg ctg att ggt gaa 834
 Ser Tyr Arg Ala Gly Tyr Leu Val Gly Asp Pro Ala Leu Ile Gly Glu
 235 240 245

35 ctc acg gaa gtc cgt aag aac ttg ggt ctc atg gtt cct ttc cca atc 882
 Leu Thr Glu Val Arg Lys Asn Leu Gly Leu Met Val Pro Phe Pro Ile
 250 255 260

40 cag cag gcc atg atc gca gcc ctc aac gac gat gac caa gag gca ggg 930
 Gln Gln Ala Met Ile Ala Ala Leu Asn Asp Asp Asp Gln Glu Ala Gly
 265 270 275 280

cag aag ctc acc tac gcg att cgt cga gca aaa ctc atg gcg gcc ctg 978
 Gln Lys Leu Thr Tyr Ala Ile Arg Arg Ala Lys Leu Met Arg Ala Leu
 285 290 295

45 ttg gaa tcc ggc ttt cag gta gat aat tct gaa gcg ggt ctg tac ctc 1026
 Leu Glu Ser Gly Phe Gln Val Asp Asn Ser Glu Ala Gly Leu Tyr Leu
 300 305 310

50 tgg gcg acg cgt gaa gaa cct tgc cgt gac act gtc gat tgg ttc gct 1074
 Trp Ala Thr Arg Glu Glu Pro Cys Arg Asp Thr Val Asp Trp Phe Ala
 315 320 325

55 gag cgt ggc att ctc gtt gcc cca gga gac ttc tat ggc cct cgc gga 1122
 Glu Arg Gly Ile Leu Val Ala Pro Gly Asp Phe Tyr Gly Pro Arg Gly
 330 335 340

60 gcg cag cat gtg cgt gtg gcg atg acc gaa acc gac gag cgc gtc gac 1170
 Ala Gln His Val Arg Val Ala Met Thr Glu Thr Asp Glu Arg Val Asp
 345 350 355 360

gcc ttt gtt tct cgc ctg agc taaacacgac taagcttatt ttgtttaatt 1221
 Ala Phe Val Ser Arg Leu Ser
 365

65 gagtttgaag ttttccgtcg aaagaggcca tttgagttcc gagtccagtc ctgagtcgag 1281

taccgagcaa aaaacctgg

1300

<210> 2
 5 <211> 367
 <212> PRT
 <213> Corynebacterium glutamicum

<400> 2
 10 Met Thr Ser Arg Thr Pro Leu Val Ser Val Leu Pro Asp Phe Pro Trp
 1 5 10 15
 Asp Ser Leu Ala Ser Ala Lys Ala Lys Ala Ala Ser His Pro Asp Gly
 20 25 30
 15 Ile Val Asn Leu Ser Val Gly Thr Pro Val Asp Pro Val Ala Pro Ser
 35 40 45
 Ile Gln Ile Ala Leu Ala Glu Ala Ala Gly Phe Ser Gly Tyr Pro Gln
 50 55 60
 Thr Ile Gly Thr Pro Glu Leu Arg Ala Ala Ile Arg Gly Ala Leu Glu
 65 70 75 80
 25 Arg Arg Tyr Asn Met Thr Lys Leu Val Asp Ala Ser Leu Leu Pro Val
 85 90 95
 Val Gly Thr Lys Glu Ala Ile Ala Leu Leu Pro Phe Ala Leu Gly Ile
 100 105 110
 30 Ser Gly Thr Val Val Ile Pro Glu Ile Ala Tyr Pro Thr Tyr Glu Val
 115 120 125
 Ala Val Val Ala Ala Gly Cys Thr Val Leu Arg Ser Asp Ser Leu Phe
 130 135 140
 Lys Leu Gly Pro Gln Ile Pro Ser Met Met Phe Ile Asn Ser Pro Ser
 145 150 155 160
 40 Asn Pro Thr Gly Lys Val Leu Gly Ile Pro His Leu Arg Lys Val Val
 165 170 175
 Lys Trp Ala Gln Glu Asn Asn Val Ile Leu Ala Ala Asp Glu Cys Tyr
 180 185 190
 45 Leu Gly Leu Gly Trp Asp Asp Glu Asn Pro Pro Ile Ser Ile Leu Asp
 195 200 205
 Pro Arg Val Cys Asp Gly Asp His Thr Asn Leu Ile Ala Ile His Ser
 210 215 220
 Leu Ser Lys Thr Ser Asn Leu Ala Ser Tyr Arg Ala Gly Tyr Leu Val
 225 230 235 240
 55 Gly Asp Pro Ala Leu Ile Gly Glu Leu Thr Glu Val Arg Lys Asn Leu
 245 250 255
 Gly Leu Met Val Pro Phe Pro Ile Gln Gln Ala Met Ile Ala Ala Leu
 260 265 270
 60 Asn Asp Asp Asp Gln Glu Ala Gly Gln Lys Leu Thr Tyr Ala Ile Arg
 275 280 285
 Arg Ala Lys Leu Met Arg Ala Leu Leu Glu Ser Gly Phe Gln Val Asp
 290 295 300

```

    Asn Ser Glu Ala Gly Leu Tyr Leu Trp Ala Thr Arg Glu Glu Pro Cys
    305                      310                      315                      320

5   Arg Asp Thr Val Asp Trp Phe Ala Glu Arg Gly Ile Leu Val Ala Pro
    325                      330                      335

    Gly Asp Phe Tyr Gly Pro Arg Gly Ala Gln His Val Arg Val Ala Met
    340                      345                      350

10  Thr Glu Thr Asp Glu Arg Val Asp Ala Phe Val Ser Arg Leu Ser
    355                      360                      365

15  <210> 3
    <211> 1300
    <212> DNA
    <213> Corynebacterium glutamicum

20  <220>
    <221> CDS
    <222> (91)..(1191)
    <223> dapC-Allele

25  <400> 3
    gggctcccca ggtggtgcgg ctaagcttgg accacaagat tttgatcacc caatgatcgc 60

    tgcgctgccg cctcaggcat aatctaacgc atg acc tct cgc acc ccg ctt gtt 114
                                1                      5
                                Met Thr Ser Arg Thr Pro Leu Val

30

    tct gtt ctt cct gat ttt ccg tgg gat tcg ctc gct tcc gca aaa gcc 162
    Ser Val Leu Pro Asp Phe Pro Trp Asp Ser Leu Ala Ser Ala Lys Ala
                                10                      15                      20

35

    aaa gct gcg tct cac ccg gat ggg atc gtg aat ctt tct gtt ggc act 210
    Lys Ala Ala Ser His Pro Asp Gly Ile Val Asn Leu Ser Val Gly Thr
    25                      30                      35                      40

40

    ccg gtt gat ccg gtc gcg ccc agc att cag atc gcg ttg gca gaa gca 258
    Pro Val Asp Pro Val Ala Pro Ser Ile Gln Ile Ala Leu Ala Glu Ala
                                45                      50                      55

45

    gcg ggg ttt tcg ggt tac cct caa acc atc ggc acc ccg gaa ctc cgc 306
    Ala Gly Phe Ser Gly Tyr Pro Gln Thr Ile Gly Thr Pro Glu Leu Arg
                                60                      65                      70

50

    gca gcc atc agg ggc gcg ctt gag cgg cgc tac aac atg aca aag ctt 354
    Ala Ala Ile Arg Gly Ala Leu Glu Arg Arg Tyr Asn Met Thr Lys Leu
                                75                      80                      85

55

    gtc gac gcc tcc ctc ctc ccc gtc gtg ggt acc aag gag gca att gcc 402
    Val Asp Ala Ser Leu Leu Pro Val Val Gly Thr Lys Glu Ala Ile Ala
                                90                      95                      100

    ctt ctt cca ttc gcg ttg ggt att tcc ggc acc gtt gtc atc cca gag 450
    Leu Leu Pro Phe Ala Leu Gly Ile Ser Gly Thr Val Val Ile Pro Glu
    105                      110                      115                      120

```

	att gcg tac cca acc tac gaa gtc gct gtc gtg gcc gca gga tgc acc	498
	Ile Ala Tyr Pro Thr Tyr Glu Val Ala Val Val Ala Ala Gly Cys Thr	
	125 130 135	
5	gtg ttg cgt tct gat tcg ctg ttt aag ctc ggc ccg cag atc ccg tcg	546
	Val Leu Arg Ser Asp Ser Leu Phe Lys Leu Gly Pro Gln Ile Pro Ser	
	140 145 150	
10	atg atg ttt atc aac tca cca tcc aac ccc aca ggc aag gtt ctg ggc	594
	Met Met Phe Ile Asn Ser Pro Ser Asn Pro Thr Gly Lys Val Leu Gly	
	155 160 165	
15	atc cca cac ttg cgc aag gtt gtg aag tgg gcg cag gaa aac aac gtg	642
	Ile Pro His Leu Arg Lys Val Val Lys Trp Ala Gln Glu Asn Asn Val	
	170 175 180	
20	atc ctc gca gct gat gaa tgc tac ttg ggt ctt ggc tgg gac gat gaa	690
	Ile Leu Ala Ala Asp Glu Cys Tyr Leu Gly Leu Gly Trp Asp Asp Glu	
	185 190 195 200	
25	aac cca ccg atc tca att ttg gat cta cgt gtc tgc gat ggc gac cac	738
	Asn Pro Pro Ile Ser Ile Leu Asp Leu Arg Val Cys Asp Gly Asp His	
	205 210 215	
30	acc aac ttg atc gcc att cac tcg ctg tct aaa acc tca aac ctc gct	786
	Thr Asn Leu Ile Ala Ile His Ser Leu Ser Lys Thr Ser Asn Leu Ala	
	220 225 230	
35	tct tac cgc gca ggt tac ctc gtt ggc gat cca gcg ctg att ggt gaa	834
	Ser Tyr Arg Ala Gly Tyr Leu Val Gly Asp Pro Ala Leu Ile Gly Glu	
	235 240 245	
40	ctc acg gaa gtc cgt aag aac ttg ggt ctc atg gtt cct ttc cca atc	882
	Leu Thr Glu Val Arg Lys Asn Leu Gly Leu Met Val Pro Phe Pro Ile	
	250 255 260	
45	cag cag gcc atg atc gca gcc ctc aac gac gat gac caa gag gca ggg	930
	Gln Gln Ala Met Ile Ala Ala Leu Asn Asp Asp Asp Gln Glu Ala Gly	
	265 270 275 280	
50	cag aag ctc acc tac gcg att cgt cga gca aaa ctc atg cgc gcc ctg	978
	Gln Lys Leu Thr Tyr Ala Ile Arg Arg Ala Lys Leu Met Arg Ala Leu	
	285 290 295	
55	ttg gaa tcc ggc ttt cag gta gat aat tct gaa gcg ggt ctg tac ctc	1026
	Leu Glu Ser Gly Phe Gln Val Asp Asn Ser Glu Ala Gly Leu Tyr Leu	
	300 305 310	
60	tgg gcg acg cgt gaa gaa cct tgc cgt gac act gtc gat tgg ttc gct	1074
	Trp Ala Thr Arg Glu Glu Pro Cys Arg Asp Thr Val Asp Trp Phe Ala	
	315 320 325	
65	gag cgt ggc att ctc gtt gcc cca gga gac ttc tat ggc cct cgc gga	1122
	Glu Arg Gly Ile Leu Val Ala Pro Gly Asp Phe Tyr Gly Pro Arg Gly	
	330 335 340	
70	gcg cag cat gtg cgt gtg gcg atg acc gaa acc gac gag cgc gtc gac	1170
	Ala Gln His Val Arg Val Ala Met Thr Glu Thr Asp Glu Arg Val Asp	
	345 350 355 360	
75	gcc ttt gtt tct cgc ctg agc taaacacgac taagcttatt ttgtttaatt	1221
	Ala Phe Val Ser Arg Leu Ser	
	365	
80	gagtttgaag ttttccgtcg aaagaggcca tttagattcc gagtccagtc ctgagtcgag	1281

taccgagcaa aaaacctgg

1300

<210> 4
 5 <211> 367
 <212> PRT
 <213> Corynebacterium glutamicum

<400> 4
 10 Met Thr Ser Arg Thr Pro Leu Val Ser Val Leu Pro Asp Phe Pro Trp
 1 5 10 15
 Asp Ser Leu Ala Ser Ala Lys Ala Lys Ala Ala Ser His Pro Asp Gly
 20 25 30
 15 Ile Val Asn Leu Ser Val Gly Thr Pro Val Asp Pro Val Ala Pro Ser
 35 40 45
 Ile Gln Ile Ala Leu Ala Glu Ala Ala Gly Phe Ser Gly Tyr Pro Gln
 50 55 60
 20 Thr Ile Gly Thr Pro Glu Leu Arg Ala Ala Ile Arg Gly Ala Leu Glu
 65 70 75 80
 25 Arg Arg Tyr Asn Met Thr Lys Leu Val Asp Ala Ser Leu Leu Pro Val
 85 90 95
 Val Gly Thr Lys Glu Ala Ile Ala Leu Leu Pro Phe Ala Leu Gly Ile
 100 105 110
 30 Ser Gly Thr Val Val Ile Pro Glu Ile Ala Tyr Pro Thr Tyr Glu Val
 115 120 125
 Ala Val Val Ala Ala Gly Cys Thr Val Leu Arg Ser Asp Ser Leu Phe
 130 135 140
 Lys Leu Gly Pro Gln Ile Pro Ser Met Met Phe Ile Asn Ser Pro Ser
 145 150 155 160
 40 Asn Pro Thr Gly Lys Val Leu Gly Ile Pro His Leu Arg Lys Val Val
 165 170 175
 Lys Trp Ala Gln Glu Asn Asn Val Ile Leu Ala Ala Asp Glu Cys Tyr
 180 185 190
 45 Leu Gly Leu Gly Trp Asp Asp Glu Asn Pro Pro Ile Ser Ile Leu Asp
 195 200 205
 Leu Arg Val Cys Asp Gly Asp His Thr Asn Leu Ile Ala Ile His Ser
 210 215 220
 Leu Ser Lys Thr Ser Asn Leu Ala Ser Tyr Arg Ala Gly Tyr Leu Val
 225 230 235 240
 55 Gly Asp Pro Ala Leu Ile Gly Glu Leu Thr Glu Val Arg Lys Asn Leu
 245 250 255
 Gly Leu Met Val Pro Phe Pro Ile Gln Gln Ala Met Ile Ala Ala Leu
 260 265 270
 60 Asn Asp Asp Asp Gln Glu Ala Gly Gln Lys Leu Thr Tyr Ala Ile Arg
 275 280 285
 Arg Ala Lys Leu Met Arg Ala Leu Leu Glu Ser Gly Phe Gln Val Asp
 290 295 300

```

      Asn Ser Glu Ala Gly Leu Tyr Leu Trp Ala Thr Arg Glu Glu Pro Cys
      305                      310                      315                      320

5     Arg Asp Thr Val Asp Trp Phe Ala Glu Arg Gly Ile Leu Val Ala Pro
      325                      330                      335

      Gly Asp Phe Tyr Gly Pro Arg Gly Ala Gln His Val Arg Val Ala Met
      340                      345                      350

10    Thr Glu Thr Asp Glu Arg Val Asp Ala Phe Val Ser Arg Leu Ser
      355                      360                      365

15    <210> 5
      <211> 32
      <212> DNA
      <213> Artificial Sequence

20    <220>
      <223> Description of Artificial Sequence: Primer

      <220>
      <223> Primer DapC (dCex1)

25    <400> 5
      gatctagaat tcgcctcagg cataatctaa cg                                     32

30    <210> 6
      <211> 32
      <212> DNA
      <213> Artificial Sequence

35    <220>
      <223> Description of Artificial Sequence: Primer

      <220>
      <223> Primer DapC (dCexna2)

40    <400> 6
      gatctatcta gacagaggac aaggcaatcg ga                                     32

```